

# **INDICATIONS OF AN INCREASE IN THE RATE OF SECTIO CAESAREA AT IHC LAVALETTE HOSPITAL**

**Jauhar Roulfia Salsabila Fatin  
Sumirah Budi Pertami, S.Kp.,M.Kep.  
Supono S.Kep, Ns, M.Kep.Sp. MB**

---

## **ABSTRACT**

Confinement sectio caesarea method is one of the alternatives for Confinement. The purpose of this study was to determine what indications are related to the increasing incidence of Sectio Caesarea at Lavalette Hospital. The type of research used in this study is descriptive research, with a cross-sectional design. The population in this study were all used as samples totaling 69 mothers giving birth. Data collection in this study used primary data by giving questionnaires to patients giving birth. Analysis was carried out using univariate, bivariate and multivariate tests. The results of the study showed that 32 (46.4%) were included in the criteria for performing a caesarean section, there was an indication of a relationship between gestational age, premature rupture of membranes (PROM), preeclampsia, breech position and prolonged labor with Sectio Caesarea procedures in mothers giving birth at Lavalette Hospital, Malang. The results of the analysis of Sectio Caesarea indications showed that C-section was the dominant indication for procedures at IHC Lavalette Hospital, Malang, with a p-value of  $0.000 < 0.005$  and the highest Odds Ratio value of 92. This means that mothers who experience have a 92-fold greater chance of undergoing a C-section compared to other variables.

**Keywords: Indications, Increase, Sectio Caesarea**