

*PICTURE OF CONSUMPTION LEVEL OF STUNTING TODDLERS BEFORE AND  
AFTER GUIDANCE IN GADING VILLAGE, MALANG REGENCY*

Lailatul Firdausi Nuzula

---

Program Studi D3 Gizi Politeknik Kesehatan Kemenkes Malang

Jl. Besar Ijen, No 77C, Kota Malang

Email : [p17110221008\\_firdausi@poltekkes-malang.ac.id](mailto:p17110221008_firdausi@poltekkes-malang.ac.id)

**ABSTRACT**

**Background:** *Baseline data results, the prevalence of stunted toddlers in Gading Village reached 20%. The purpose of this study was to determine the description of assistance related to the level of consumption of stunted toddlers in Gading Village, Malang Regency. **Research Methods:** This type of research is descriptive research with a case study design. This study was conducted on August 9 - August 28, 2024 on toddlers who were indicated as stunting. The subjects taken were 2 stunted toddlers. **Results of the study:** Nutrition assistance in the form of counseling for 4 times in 4 weeks can increase the level of toddler consumption, especially in energy and protein consumption. **Conclusion:** The percentage of energy adequacy in the first toddler is included in the category according to the AKG, but protein, carbohydrates, Fe, and Vitamin D are in the category of less, while fat, vitamin A and zinc are in the category of more. The percentage of energy adequacy in the second toddler is included in the category of more than the AKG, protein and carbohydrates are included in the category according to the AKG, but fat, vitamin A, and Zn are included in the category of more, and Vitamin D is in the category of very less.*

**Keywords:** *Stunting, Nutrition Assistance, Conseling, Consumption Level.*